

SUSTOL[®] (granisetron) extended-release injection

Coding Reference Guide

ICD-10 Diagnosis Code

Code	Description
R11.0	Nausea
R11.10	Vomiting, unspecified
R11.11	Vomiting without nausea
R11.12	Projectile vomiting
R11.13	Vomiting of fecal matter
R11.14	Bilious vomiting
R11.2	Nausea with vomiting, unspecified
T45.1X5*	Adverse effects of antineoplastic and immunosuppressive drugs
Z41.9*	Encounter for other procedures for purposes other than remedying health state
Z51.11**	Encounter for antineoplastic chemotherapy

*Supplementary Classification Code

**Required when given within 48 hours of moderately or highly emetogenic chemotherapy

National Drug Code (NDC)

NDC	Description
47426-0101-06	Single-dose kit, 10 mg granisetron/0.4 mL

Coding for SUSTOL

HCPCS Code	Description
J1627	Injection, granisetron extended-release, 0.1 mg

Professional Services and CPT Codes

CPT Code	Description
96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular

Hospital Service and Supplies

Revenue Codes	Description
0631	Single source drug
0636	Drugs requiring detailed coding

The coding information contained herein is for informative purposes only, and is not a guarantee of coverage or reimbursement for any product or service. This information is not intended to substitute for the physician's independent diagnosis or treatment of each patient. Coding requirements may vary by payer; please consult the payer to determine which codes are required.

For questions regarding SUSTOL billing and coding please call Heron Connect at **1-844-HERON11 (1-844-437-6611)** from 8 AM to 8 PM ET, Monday through Friday.

Please see Indications and Important Safety Information on reverse.

www.sustol.com



sustol[®]
(granisetron) extended-release injection

SUSTOL® (granisetron) extended-release injection

Indication¹

SUSTOL is indicated in combination with other antiemetics in adults for the prevention of acute and delayed nausea and vomiting associated with initial and repeat courses of moderately emetogenic chemotherapy (MEC) or anthracycline and cyclophosphamide (AC) combination chemotherapy regimens.

Important Safety Information

Contraindications

SUSTOL is contraindicated in patients with hypersensitivity to granisetron, any of the components of SUSTOL, or any other 5-HT₃ receptor antagonist.

Warnings and Precautions

Injection site reactions (ISRs), including infection, bleeding, pain and tenderness, nodules, swelling, and induration, have occurred with SUSTOL. Monitor for ISRs following SUSTOL injection. Inform patients that some ISRs may occur 2 weeks or more after SUSTOL administration. In patients receiving antiplatelet agents or anticoagulants, consider the increased risk of bruising or severe hematoma prior to the use of SUSTOL.

Monitor for constipation and decreased bowel activity and consider optimizing patients' current bowel regimens used for managing preexisting constipation. Instruct patients to seek immediate medical care if signs and symptoms of ileus occur.

Hypersensitivity reactions have been reported and may occur up to 7 days or longer following SUSTOL administration and may have an extended course. If a reaction occurs, administer appropriate treatment and monitor until signs and symptoms resolve.

Serotonin syndrome has been reported with 5-HT₃ receptor antagonists alone but particularly with concomitant use of serotonergic drugs.

Use in Specific Populations

Avoid SUSTOL in patients with severe renal impairment. In patients with moderate renal impairment, administer SUSTOL not more frequently than once every 14 days.

Adverse Reactions

Most common adverse reactions ($\geq 3\%$) are injection site reactions, constipation, fatigue, headache, diarrhea, abdominal pain, insomnia, dyspepsia, dizziness, asthenia, and gastroesophageal reflux.

For more information about SUSTOL, please see accompanying full Prescribing Information.

Reference: 1. SUSTOL [package insert]. Heron Therapeutics, Inc., San Diego, CA; May 2017.